



## The Study of Multidrug Resistance Escherichia Coli in Drinking Water Samples from 9<sup>th</sup> District of Kabul City

Dr.Ahmad Wali Ataye<sup>1</sup>, Dr.Ehsanullah Ehsan<sup>2</sup>, Dr. Farida Razavi<sup>3</sup>, Dr. Abdul Moaid Kohistani<sup>4</sup>, Qudratullah Ebrahimi<sup>5</sup>, Noorullah Mandeza<sup>6</sup>

<sup>1</sup>MD, MPH and DCH, Microbiology Department, Kabul University of Medical Science “Abu Ali Ibn Sina”, Kabul, Afghanistan, <sup>2</sup>MD, Dermatology Department, Alberoni University, <sup>3</sup>MD, Pediatric surgery Department, Kabul University of Medical Science “Abu Ali Ibn Sina”, Kabul, Afghanistan, <sup>4</sup>MD, Internal medicine Department, Alberoni University, <sup>5</sup>Qudratullah Ibrahim, BSc of Medical Laboratory Technology Kabul University of Medical Science “Abu Ali Ibn Sina”, Kabul, Afghanistan, <sup>6</sup>Faculty member of Biochemistry department at Dawat university

drataye1@gmail.com

**Abstract.** Antibiotic resistance in pathogenic Bacteria is a serious public health issue. Various factors contribute to the spread of antibiotic –resistance microorganisms in the environment. This study was aimed for detection of multi drug resistance Escherichia Coli in drinking water samples from 9<sup>th</sup> district of Kabul city..Water sample is taken in Simple Random Sampling. The population size of drinking water sampling is N=378 and by using of Epi – info program, the water sample is selected and so, the sample size of water sample is 378 samples, therefore 126 water sample from house hold ,126 samples from supplies and 126 samples from collection tankers of 9<sup>th</sup> district of Kabul city. Samples of drinking water from district 9<sup>th</sup> of Kabul city were collected in a sterile bottle from various parts of district 9th of Kabul city bringing into Microbiology Department and growing it in mac-conkey agar. the collected data is by using of MS –excel and SPSS software was analyzed. out of 378 samples collected in 243 (64.3%) Escherichia coli were isolated and 135(35.7%) samples were free of E.coli. Maximum resistance was observed against Nalidixic acid( 64.3%) followed by ampicilline) 59.5%),Ciprofloxacin,( 57.1%.It is also remarkable that percentage of Escherichia coli in different sources of 9<sup>th</sup> district of Kabul city such as house hold, supplies and collection tanker was as follows: 27.1%,30.8% and 41.9%. In conclusion this study found that contaminated percentage of drinking water in 9<sup>th</sup> district of Kabul city was higher 64.3% and presence of E-coli is higher in tanker collection and antibiotic resistance is higher against Nalidixic acid .Further studies is needed for better results in future.

**Keywords.** Drinking water, Escherichia coli, antibiotic resistance, multidrug resistance, 9th district of Kabul city

### 1.Introduction:

Antibiotic resistance in pathogenic Bacteria is a serious public health problem. Various factors contribute to the spread of antibiotic–resistance microorganisms in the environment. Faecal coli forms are a group of bacteria, which are natural inhabitants of the gut of humans

and other warm-blooded animals. Escherichia Coli (E. coli) is a member of fecal coli forms that contaminate the drinking water from human and animal fecal waste. During rainfalls, these coli forms may be washed into creeks, rivers, streams, lakes, or ground water. Untreated drinking water coming from these sources contains coliforms including E. coli. In developing countries, drinking water supply lines and open sewage drains are laid side by side resulting in frequent contamination of water.<sup>1</sup> E.coli is an opportunistic pathogen in neonatal and immunocompromised patients.<sup>2</sup> Bacteremia, wound infections, urinary tract infections, and gastrointestinal infections are the diseases associated with E.coli and are often fatal in newborns.<sup>3</sup> Food and water borne outbreaks of E. coli have been documented from a number of countries.<sup>4,5</sup> The difficulties in the treatment of food and water associated gastrointestinal diseases due to E. coli have been reported. This problem is compounded by the continued emergence of antibiotic resistance to a growing number of antibiotics; i.e. Carbenicillin, Tetracycline, Streptomycin,<sup>6</sup> Norfloxacin, Amoxycillin, Trimethoprim, Nitrofurantoin,<sup>7</sup> Nalidixic acid, Gentamicin, Cefuroxime,<sup>8</sup> etc. Increase in antibiotic resistance level is now a global problem. Infections with antibiotic resistant bacteria makes the therapeutic options for infection treatment, extremely difficult or virtually impossible in some instances.<sup>9</sup> Therefore, the determination of antimicrobial susceptibility of clinical isolate is often crucial for optimal antimicrobial therapy of infected patients. A high-density patient population in frequent contact with health care staff and the attendant risk of cross-infection contributes to the spread of antibiotic-resistant micro-organisms in the environment.<sup>10</sup> Occurrence and prevalence of these resistant strains in environment is therefore, a usual kind of thing in the developing countries. Since water is one of the four components of environment, and a usual habitat for E. coli, therefore, the availability of antibiotic resistant E. coli strains in water cannot be denied. The multi-drug resistant (MDR) E. coli strains in drinking water have been reported by Walia SK.<sup>6</sup> Our study is aimed for detection of multi drug resistance Escherichia Coli in drinking water samples (household, supplies and collection tanker) of district 9<sup>th</sup> of Kabul city.

### **Review Literatures:**

Escherichia coli is normal flora of human gastrointestinal system and is isolated from stool of humans.<sup>15</sup> Escherichia coli is one of the most common bacteria which cause diarrhea in humans. Pathogenic species of Escherichia Coli which are included enter pathogenic Escherichia coli, enterotoxigenic Escherichia coli, enteroinvasive Escherichia coli and enteroaggregative Escherichia coli, they could cause diarrhea.<sup>16</sup> enterotoxigenic Escherichia coli are present in food and water resources of developing countries and it also caused diarrhea among children under age of two years in developing countries. Approximately a healthy person swallows 10<sup>8</sup> up to 10<sup>10</sup> enterotoxigenic Escherichia coli, it could cause diarrhea.<sup>17</sup> Also enterotoxigenic Escherichia coli among those passengers which are traveling to tropical and subtropical countries and faced into contaminated food and water, it may cause diarrhea. Moreover, enterotoxigenic Escherichia coli also may be one of bacterial cause of diarrhea in developed countries of the world.<sup>18,19</sup> Escherichia coli could easily grow under aerobic conditions, and on mac-conkey agar and could produce pink colony and it shows that Escherichia coli ferments glucose. The most significant chemical reaction for identification of Escherichia coli is indole test and at 99% it is indole positive.<sup>20</sup> A lot of research studies have been conducted to identify the multidrug resistance Escherichia coli in drinking water. One study which has been conducted in drinking water of Alhasa region of Saudi Arabia in 2010, it shows that contamination percentage of drinking water with Escherichia coli is 86.7%. As 57.7% of isolated Escherichia coli is at least resistant against three antimicrobial drugs such as

Ampicillin, Nalidixic acid and Ciprofloxacin and 42.3% of isolated Escherichia coli is resistant against one or more than one antimicrobial drugs.<sup>21</sup> Moreover, a study in Meso state of India in 2016, shows that 54% of isolated Escherichia coli in drinking water is resistant against antimicrobial drugs, 16% of isolated Escherichia coli in drinking water has intermediate effect against antimicrobial drugs and 30% of isolated Escherichia coli is sensitive against antimicrobial drugs.<sup>22</sup> Problem statement, Questions and Tentative Hypothesis Water is a basic requirement of life and health and a basic nutrient of the human body. Water related diseases continue to be one of the major health problems globally. Everyday water related diseases cause the death of thousands of children, untold sufferings and loss of working time. About 4 billion cases of diarrhea occur annually worldwide of which 88% is attributed to unsafe water supply, inadequate sanitation and hygiene. Kabul city is the capital and most populated city in the country. The estimated population of Kabul city is approximately five million. This city has been facing different challenges like overgrowing population and rapid urbanization. Situations of water and environmental sanitation and hygiene have deteriorated in the city; Kabul city was initially designed for the population of one million but is now home to population around four million.

### **Hypothesis**

The Contamination level with multi- drug resistant E. Coli is high among different sources of water in Kabul city.

### **Research Questions:**

- 1) What is the percentage of Multi drug resistance Escherichia Coli in drinking water of District 9<sup>th</sup> of Kabul city?
- 2) The Multi drug resistance Escherichia Coli is resistant against which antimicrobial drugs?

### **Research Objectives:**

- 1) This study is aimed to detect multi drug resistance Escherichia Coli in drinking water samples (household, supplies and collection tanker) of district 9<sup>th</sup> in Kabul city.
- 2) To identify whether there exists any difference regarding water contamination of district 9<sup>th</sup> of Kabul city and different sources of water.

### **Inclusion criteria:**

Drinking water of household, water supply and collection tanker of district 9<sup>th</sup> of Kabul city.

### **Exclusion criteria :**

Drinking water of other sources of other districts of Kabul city.

## **2.Methods and materials:**

This study is a descriptive Cross Sectional done in spring of 2021( March –April-May) . Water samples were taken using Simple Random Sampling method from District 9<sup>th</sup> of Kabul city were selected by using the convenient sampling method, from water sources of household ,suppliers and from collection tankers.

The population size of drinking water sample is N=378 and by using of Epi – info program, the water sample is selected and so, the sample size of water sample is 378 samples,

therefore 126 water sample from house hold ,126 samples from suppliers and 126 samples from collection tankers of 9<sup>th</sup> district of Kabul city

Samples of drinking water will be collected in a sterile bottle from various locations of 9th district of Kabul city. Immediately the water sample would be brought to Microbiology Department and growing it in mac-conkey agar. The collected data would be analyzed by using of Ms Excel and SPSS software .

### 3.Results :

Out of 378 samples collected in 243 (64.3%) Escherichia coli were isolated and 135(35.7%) samples were free of E.coli. Maximum resistance was observed against Nalidixic acid (64.3%) followed by ampicilline (59.5%), Ciprofloxacin, (57.1% ( and ceftriaxone (59.5%(. It is also remarkable that percentage of Escherichia coli in different sources of 9th district of Kabul city such as house hold, supplies and collection tanker was as follows:27.1%,30.8% and 41.9% . in this study it was found, that contaminated percentage of drinking water in 9 th district of Kabul city was higher .Tables(2,3,4).

### 4.Discussions:

Since their discovery, antimicrobial drugs have been proved remarkably effective for the control of bacterial infections. However, it was soon evidenced that bacterial pathogens were unlikely to surrender unconditionally, and some pathogens rapidly became resistant to many of the first effective drugs<sup>23</sup>.

Over the past few decades, antimicrobials have become increasingly available for a broad range of pathogens. Due to the widespread use of these drugs, new forms of antimicrobial resistance have emerged <sup>24</sup>. Antibiotic resistance profile for clinical E. coli isolates have well been documented by various workers. Opportunistic pathogens presenting broad-spectrum antibiotic resistance have emerged extensively in hospital environments, causing serious infections in immunocompromised hosts<sup>25</sup>. However, less attention is given to environmental E. coli isolates. Almost same results for E. coli strains isolated from drinking water have been reported by Walia SK in 2004 .

### 5.Conclusion:

In this study it was found that contaminated percentage of drinking water in 9<sup>th</sup> district of Kabul city was higher 64.3% and presence of E-coli is higher in tanker collection and antibiotic resistance is higher against Nalidix acid .Further studies is needed for better results in future.

Table(1).show the different sources of drinking water in district 9<sup>th</sup> of Kabul city.

Water source	Frequency	Percentage	Total percentage
House hold	126	33.3 %	33.3 %
Supplies	126	33.3 %	33.3 %
Collection Tankers	126	33.3 %	33.3 %
Total	378	100 %	100%

Table 2: Show the percentage of E.coli in drinking Water of district 9th of Kabul city

The presence of E.coli in drinking water	Frequency	Percentage	Total percentage
Yes	243	64.3%	64.3%
No	135	35.7%	35.7%
Total	378	100%	100%

Table 3: Show the percentage of E.coli in different sources drinking water of 9<sup>th</sup> district of Kabul city

The percentage of E.coli in house hold source	27.1% (22samples)
The percentage of E.coli in supplies sources	30.87% (25samples)
The percentage of E.coli in collection tanker	41.98% (34 samples)

Table 4 Show the Result of Antibigram of different Antibiotics for drinking water of 9<sup>th</sup> district in Kabul city.

Antibiotics	Frequency	Percentage
Nalidixic acid Resistance	81	64.3%
Ampicilline Sensitive Resistance	79	59.5%
Ceftriaxone Sensitive Resistance	79	59.5%
Ciprofloxacin Sensitive Resistance	78	57.1%

### 6.Conflict of interest:

The authors have no conflict of interest.

### References

- [1] Ahmed, I., et al.(2010). Hepatitis E in Pakistan.East Mediterr Health J.pp. 121-128 .
- [2] Alrashed, K.M.,et al.(2011). Resistant gram-negative bacilli and antibiotic consumption in Zarqa, Jordan. Pak J Med Sci.pp; 63-59 .
- [3] Bax,R.,et al.(2002).The future challenges facing the development of new antimicrobial drugs. Nat Rev Drug Discovery.pp. 910-895 .Retrieved from

<http://www.ncbi.nlm.nih.gov/pubmed>.

[4] Becker, S., et al. (1982). Contamination of weaning foods and transmission of enterotoxigenic *Escherichia coli* diarrhoea in children in rural Bangladesh. *Trans R Soc Trop Med Hyg*. Pp. 276-259 .Retrieved from <http://WWW.Uptodate.com/Contents/abstract>.

[5] Chaudhry, G.R., et al. (2004). Self-transmissible Antibiotic Resistance to Ampicillin, Streptomycin, and Tetracycline found in *Escherichia coli*. Isolates from contaminated Drinking water. *J Environmental Sci and Health*. pp. 662- 651.

[6] Carey, R.B., et al. (2011). Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria. An international expert proposal for interim standard definitions for acquired resistance. *Clinical Microbiology Infection*. Pp. 1491-1461 .Retrieved from

<http://www.ncbi.nlm.nih.gov/pubmed>.

[7] Coates, A., et al. (2002). The future challenges facing the development of new antimicrobial drugs. *Nat Rev Drug Discovery*. pp. 910-895 .Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed>.

[8] WWW, (2011). [cdc.gov](http://www.cdc.gov).

[9] El-Astal, Z. (2010). Bacterial pathogens and their antimicrobial Susceptibility in Gaza Strip, Palestine. *Pak J Med Sci*. pp. 365-70.

[10] Guerra, B., et al. (2006). Phenotypic and genotypic characterization of antimicrobial resistance

in *Escherichia coli* O111 isolates. *J Antimicrob Chemother*. pp. 1214-1210 . Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed>.

[11] Howard, D.H., et al. (2003). The Global Impact of Drug Resistance. *Clinical Infectious Disease*. p. 36. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed>.

[12] Khan, M.H., et al. (2011). Antibacterial activity of natural spices on multiple drug resistant *Escherichia coli* isolated from drinking water. Bangladesh. *Ann Clinical Microbiology Antimicrobiology*. p. 10. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed>.

[13] Kibret, M., et al. (2011). Antimicrobial susceptibility patterns of *E. coli* from clinical sources in northeast Ethiopia. *Afr Health Sci*. pp. 45-40 .Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed>.

[14] Kumar, A.K., et al. (2004). Prevalence of extended-spectrum  $\beta$ -lactamase among multidrug resistant Gram-negative isolates from a general hospital in Saudi Arabia. *Saudi Med J*. pp. 574-570 .Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed>.

[15] Kaper, J.B., et al. (1998). Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev* .pp. 142-11 .Retrieved from <http://WWW.Uptodate.com/Contents/abstract>.

[16] Lewis, D., et al. (2010). The fate of *Escherichia coli* O157 in soil and its potential to contaminate drinking water. *Int J Food Microbiology*. pp. 117-111 .

[17] Monica, Cheesbrough, (2000). *District Laboratory Practice in Tropical Countries*. p. 137.

[18] Pyrrho, A.S., et al. (1999). Antibody response in mice immunized with a plasmid DNA encoding the colonization factor antigen I of enterotoxigenic *Escherichia coli*. *FEMS Immunol Med Microbiol* .pp. 321-323 . Rev .pp. 142-11 -142 .Retrieved from <http://WWW.Uptodate.com/Contents/abstract>.

[19] Pierce, N.F., et al. (1973). Differences in the response of rabbit small intestine to heat-labile and heat-stable enterotoxins of *Escherichia coli*. *Infect Immun*. pp. 142- 11. Retrieved from <http://WWW.Uptodate.com/Contents/abstract>.

[20] Osterblad, M., et al. (2000). A between-Species Comparison of Antimicrobial Resistance in Enterobacteria in Fecal Flora. *Antimicrobia Agents and Chemotherapy* .pp. 1484-1478 .



- [21] Rowe ,B., et al.(1981). Enterotoxigenic Escherichia coli diarrhoea: acquired immunity and transmission in an endemic area. *Bull World Health Organ*.pp. 142-111.Retrievedfrom<http://WWW.Uptodate.com/Contents/abstract>.
- [22] Rothbaum, R.,et al.(1982). A clinicopathologic study of enterocyte-adherent Escherichia coli: a cause of protracted diarrhea in infants. *Gastroenterology*.pp. 441- 483. Retrieved from <http://WWW.Uptodate.com/Contents/abstract>.
- [23] Ratnam, S.,et al.(1986). Sorbitol-MacConkey medium for detection of Escherichia coli O157:H7 associated with hemorrhagic colitis. *J Clinical Microbiology*.pp. 142-111 .Retrievedfrom<http://WWW.Uptodate.com/Contents/abstract>.
- [24] Ulshen ,M.H.,et al.(1980). Pathogenesis of Escherichia coli gastroenteritis in man-anothermechanism.*N Engl J Med*.Pp.302-99.Retrievedfrom<http://WWW.Uptodate.com/Contents/abstract>.
- [25] Sommerfelt, H.,et al.(1989). Enteroaggregative Escherichia coli and Salmonella associatedwithnondysentericpersistentdiarrhea.
- [26] Sorum, H.,et al. (1994). Transfer of multiple drug resistance plasmids between bacteria of diverse origins in natural microenvironments. *Appl. Environ.Microbiology*.pp. 1415-1421.